

# Jon the Quantum Mechanic

Chris Quigg

*Fermi National Accelerator Laboratory*



Jon Rosner Symposium · University of Chicago · 1 April 2011

# First DPF Meeting · Boulder

Program for the Discussion Session On

## Quark Models

Discussion Leader:  
P. Freund

Date and time: 2:15-5:15 p.m., August 19, 1969  
Place: West Ballroom

P. G. O. Freund - Hadron Dynamics and Quarks

J. L. Rosner - Quarks and Crossing Symmetry



G. Zweig - A Model for the Hadrons

R. H. Capps - Baryonic Exchange Degeneracies

H. J. Lipkin - Algebraic Structure and Spin Couplings  
In the Levin-Frankfurt Model  
(No paper submitted)

S. Meshkov - Experimental Consequences of  $SU(6)_W$

# June 16, 1977: News of $\Upsilon$



Quarkonium? One new quark or two?  
Properties of new quark?  
Is strong interaction flavor-blind?

# The Empirical Approach

- Appelquist & Politzer: NRQM applies to  $Q\bar{Q}$  systems.
- Cornell group: had shown the predictive power of the NR potential-model approach using a “culturally determined” potential,

$$V(r) = -\frac{\kappa}{r} + \frac{r}{a^2} .$$

- Eichten & Gottfried anticipated spectroscopy of  $b\bar{b}$ :

$$\begin{aligned} M(\Upsilon') - M(\Upsilon) &\approx 420 \text{ MeV} \\ &\approx \frac{2}{3}[M(\psi') - M(\psi)] . \end{aligned}$$

$\Upsilon' - \Upsilon$  spacing same as  $\psi' - \psi$

E288	$M(\Upsilon') - M(\Upsilon)$	$M(\Upsilon'') - M(\Upsilon)$
Two-level fit	$650 \pm 30$ MeV	
Three-level fit	$610 \pm 40$ MeV	$1000 \pm 120$ MeV
$M(\psi') - M(\psi)$	$\approx 590$ MeV	

$V(r) = C \log r \rightsquigarrow \Delta E$  independent of  $\mu$

# Scaling the Schrödinger Equation: $V(r) = \lambda r^\nu$

$$\frac{\hbar^2}{2\mu} u'' + \left[ E - \lambda r^\nu - \frac{\ell(\ell+1)\hbar^2}{2\mu r^2} \right] u(r) = 0 .$$

Substitute

$$r = \rho \left( \frac{\hbar^2}{2\mu|\lambda|} \right)^{1/(2+\nu)}$$

$$E = \varepsilon \left( \frac{\hbar^2}{2\mu|\lambda|} \right)^{-2/(2+\nu)} \left( \frac{\hbar^2}{2\mu} \right)$$

and identify  $w(\rho) \equiv u(r)$ :  $\leadsto$  dimensionless form,

$$w''(\rho) + \left[ \varepsilon - \text{sgn}(\lambda) \rho^\nu - \frac{\ell(\ell+1)}{\rho^2} \right] w(\rho) = 0 .$$

Length scale [ $L$ ]  $\propto (\mu|\lambda|)^{-1/(2+\nu)}$

Coulomb:	$(\mu \lambda )^{-1}$
Log: $V(r) = C \log r$	$(C\mu)^{-1/2}$
Linear:	$(\mu \lambda )^{-1/3}$
SHO:	$(\mu \lambda )^{-1/4}$
Square well:	$(\mu \lambda )^0$

Energies [ $\Delta E$ ]  $\propto (\mu)^{-\nu/(2+\nu)}(|\lambda|)^{2/(2+\nu)}$

Coulomb:	$\mu \lambda ^2$
Log: $V(r) = C \log r$	$C\mu^0$
Linear:	$\mu^{-1/3} \lambda ^{2/3}$
SHO:	$\mu^{-1/2} \lambda ^{1/2}$
Square well:	$\mu^{-1}$

# Measuring the $b$ -quark's charge

For  $\nu \leq 1$ , scaling laws imply


$$|\Psi_b(0)|^2 \geq \frac{m_b}{m_c} |\Psi_c(0)|^2$$

$$\leadsto \Gamma(\Upsilon_n \rightarrow \ell^+ \ell^-) \geq \frac{e_b^2}{e_c^2} \cdot \frac{m_b}{m_c} \cdot \frac{M(\psi_n)^2}{M(\Upsilon_n)^2} \Gamma(\psi_n \rightarrow \ell^+ \ell^-)$$

DORIS results presented at 1978 ICHEP (Tokyo)

$$\begin{aligned}\Gamma(\Upsilon \rightarrow \ell^+ \ell^-) & 1.26 \pm 0.21 \text{ keV} \\ \Gamma(\Upsilon' \rightarrow \ell^+ \ell^-) & 0.36 \pm 0.09 \text{ keV}\end{aligned}$$


established  $e_b = -\frac{1}{3}$



# Counting Narrow Levels of Quarkonium

Eichten–Gottfried:  $Q\bar{Q}$  ( $m_Q \gg m_c$ ):  $\geq 3$  narrow  ${}^3S_1$  levels, but  $\Upsilon' - \Upsilon$  spacing is much larger than predicted.

How general? Depends on?



Logarithmic potential: 3 or 4 narrow  $\Upsilon({}^3S_1)$  levels.

# Counting Narrow Levels . . .

Remarkable general result

Number of narrow  ${}^3S_1$  levels is




$$n \approx 2 \cdot \left( \frac{m_Q}{m_c} \right)^{1/2}$$

Key ingredients:

- $\delta(m_Q) \equiv 2m(\text{lowest } Q\bar{q} \text{ state}) - 2m_Q$  approaches a finite  $\delta_\infty$  independent of  $m_Q$ .
- Semiclassical (WKB) approximation:

$$\int_0^{r_0} dr [m_Q(\delta(m_Q) - V(r))]^{1/2} = (n - \frac{1}{4})\pi$$

# How $n \propto \sqrt{m_Q}$ is realized



# Semiclassical Methods and Results

Evaluating the nonrelativistic connection

$$|\Psi_n(0)|^2 = \frac{\mu}{2\pi} \left\langle \frac{dV}{dr} \right\rangle_n$$

in semiclassical approximation, connect the square of the s-wave wave function at the origin to the level density:

$$|\Psi_n(0)|^2 = \frac{(2\mu)^{3/2}}{4\pi^2} E_n^{1/2} \frac{dE_n}{dn}$$

(for a nonsingular potential).

## Elementary application

For  $V(r) = \lambda r$ ,  $|\Psi_n(0)|^2$ : independent of  $n$ ,  $\leadsto E_n \sim n^{2/3}$ .

# Semiclassical Inverse Problem


For a monotonically increasing potential, the semiclassical quantization condition

$$\int_0^{r_0} dr \{2\mu[E - V(r)]\}^{1/2} = (n - \frac{1}{4})\pi$$

connects the shape of the potential to the level density:

$$r(V) = \frac{2}{2\mu^{1/2}} \int_0^V dE (V - E)^{1/2} \left[ \frac{dE_n}{dn} \right]^{-1} .$$

New results, or results forgotten for 40 years?



ISIDOR I. RABI  
850 RIVERSIDE DRIVE  
NEW YORK, NEW YORK 10027.

Dear Dr Quigg,

I don't remember very  
clearly who first did


$$|\psi(0)|^2 = \frac{u}{2\pi} \left(\frac{dV}{dx}\right). I$$

always thought Fermi.  
I suggest you look at  
some of the early papers  
on HFS about 1931 for references.

Of course Casimir did  
the real job and you will  
find full references there.

I am sure I cannot  
reply more fully but Casimir  
will do the job. Legre will  
also know. Sincerely yours

J.J. Rabi



UNIVERSITY OF CALIFORNIA, BERKELEY

BERKELEY • DAVIS • DIVINE • LOS ANGELES • RIVERSIDE • SAN DIEGO • SAN FRANCISCO

DEPARTMENT OF PHYSICS



SANTA BARBARA • SANTA CRUZ

BERKELEY, CALIFORNIA 94720

MARCH 30 1979

DR. C. QUIGG  
Fermilab  
Batavia IL

Dear Dr. Quigg:

in reply to your letter of March 19th. I do not remember having seen the formula for  $|\psi(0)|^i$  that you quote. Fermi has had repeated occasions for calculating  $|\psi(0)|$  and you may find it in papers # 75b, 95 of the Collected papers. It was a quantity very familiar to him and on which he returned often. However I do not recall the specific formula you quote. In what context did you obtain it?

Sincerely yours

Emilio Segrè

ISSN 0370-1573

# PHYSICS REPORTS

A Review Section of Physics Letters

## QUANTUM MECHANICS WITH APPLICATIONS TO QUARKONIUM

C. QUIGG and J.L. ROSNER

Volume 56 Number 4

December 1979

PRPLCM 56(4) 167-235 (1979)



NORTH-HOLLAND PUBLISHING COMPANY  
AMSTERDAM

## Dualities

Connect bound-state spectra of  $V(r) = \lambda r^\nu$  ( $\nu > 0$ )  
and  $\bar{V}(r) = \bar{\lambda} r^{\bar{\nu}}$  ( $\bar{\nu} < 0$ )

### Paired Schrödinger equations


$$\frac{\hbar^2}{2\mu} u''(r) + \left[ E - \lambda r^\nu - \frac{\ell(\ell+1)\hbar^2}{2\mu r^2} \right] u(r) = 0$$

$$\frac{\hbar^2}{2\mu} v''(z) + \left[ \bar{E} - \bar{\lambda} z^{\bar{\nu}} - \frac{\bar{\ell}(\bar{\ell}+1)\hbar^2}{2\mu z^2} \right] v(z) = 0$$


$$(\nu + 2)(\bar{\nu} + 2) = 4, \quad \bar{E} = \lambda (\bar{\nu}/\nu)^2, \quad \bar{\lambda} = -E (\bar{\nu}/\nu)^2,$$
$$z = r^{1+\nu/2}, \quad (\bar{\ell} + 1/2)^2 \nu^2 = (\ell + 1/2)^2 \bar{\nu}^2$$

familiar case Coulomb  $\iff$  harmonic oscillator

# Our Priority Dispute with Isaac Newton



S. Chandrasekhar



# **Classical orbits in power-law potentials**

Aaron K. Grant and Jonathan L. Rosner

*Enrico Fermi Institute and Department of Physics, University of Chicago, Chicago, Illinois 60637*

(Received 10 May 1993; accepted 11 November 1993)

The motion of bodies in power-law potentials of the form  $V(r) = \lambda r^\alpha$  has been of interest ever since the time of Newton and Hooke. Aspects of the relation between powers  $\alpha$  and  $\bar{\alpha}$ , where  $(\alpha+2)(\bar{\alpha}+2)=4$ , are derived for classical motion and the relation to the quantum-mechanical problem is given. An improvement on a previous expression for the WKB quantization condition for nonzero orbital angular momenta is obtained. Relations with previous treatments, such as those of Newton, Bertrand, Bohlin, Fauré, and Arnold, are noted, and a brief survey of the literature on the problem over more than three centuries is given.

Am. J. Phys. 62, 310 (1994)

## Designer Potentials

Hank Thacker

Construct a symmetric, one-dimensional potential that supports  $N$  bound states at specified  $E_n$  out of reflectionless potentials

$$V(x) = -2\kappa^2 \operatorname{sech}^2[\kappa(x - x_0)]$$


(single bound state at  $E = -\kappa^2$ )

$N$ -level reflectionless potential:  
 $N$ -solitary-wave solution to



$$v_t - 6vv_x + v_{xxx} = 0$$

Korteweg-de Vries equation

# Reflectionless potentials as Korteweg-de Vries solitons: harmonic oscillator example




# Flavor independence of strong interaction among heavy quarks



Reconstructed from  $\Psi$


Reconstructed from  $\Upsilon$

# No Degenerate Levels in One-Dimensional QM (simple Wronskian proof, if no pathologies)




*As levels approach, two buckets retreat to  $\pm\infty$*


# Band Structure and Periodic Potentials




# Band Structure and Periodic Potentials




# Charmonium (PDG)




# BaBar $\eta_c(1S)$ , $\chi_{c0}(1P)$ , $\chi_{c2}(1P)$ , $\eta_c(2S)$ in $\gamma\gamma$




# $B_c$ (6277)




# Bottomonium (PDG)



# BaBar $h_b(\text{1P})$ : $\Upsilon(3\text{S}) \rightarrow \pi^0 h_b(\text{1P}) \rightarrow \gamma \eta_b(\text{1S})$



# BELLE $h_b(1P)$ and $h_b(2P)$ : $\pi^+\pi^- + \text{MM}$



# New States Associated with Charmonium

- $X(3872) \rightarrow \pi^+ \pi^- J/\psi$  at  $D^0 \bar{D}^{*0}$  threshold;  
very likely  $J^{PC} = 1^{++}$ ;  
 $(c\bar{c}) + s\text{-wave threshold} + \text{"molecule"} + \dots?$   
Feshbach resonances?      coupled-channels
- $Z(3930)$  in  $\gamma\gamma \rightarrow D\bar{D}$ :  $\chi'_{c2}$
- $X(3940)$  in  $e^+ e^- \rightarrow J/\psi D^* \bar{D}$
- $Y(3940)$  in  $B \rightarrow K\omega J/\psi$
- $X(3915)$  in  $\gamma\gamma \rightarrow \omega J/\psi \sim Z(3930)?$
- $Y(4140)$  in  $J/\psi \phi$
- $Y(4260)$  in  $e^+ e^- \rightarrow \gamma_{\text{ISR}} J/\psi \pi^+ \pi^-$
- $Z^+(4430) \rightarrow \pi^+ \psi'$ : not yet confirmed  
 $2^{--}, 2^{-+}$  states (above  $D\bar{D}$ , but narrow) still missing

富士山

富士山

